

**Sutter Medical Center Santa Rosa
Antibiogram 2012**

Organism Name	CountOfOrg code	Amikacin	Ampicillin	Ampicillin/Sulbactam	Cefazolin	Cefepime	Cefoxitin	Ceftazidime	Ceftriaxone	Ciprofloxacin	Gentamicin	Imipenem	Levofloxacin	Nitrofurantoin***	Tobramycin	Trimethoprim/Sulfa	Piperacillin/Tazobactam Count	Piperacillin/Tazobactam
Enterobacter cloacae⁺	22	100%				100%		91%	83%	96%	96%	100%	96%	31%	96%	83%		
Escherichia coli⁺⁺	520	99%	57%	70%	91%	97%	93%	96%	96%	88%	93%	99%	88%	94%	93%	76%	(78)	94%
Klebsiella oxytoca⁺	10	100%		73%	64%	91%	80%	91%	91%	91%	91%	100%	91%	86%	100%	82%		
Klebsiella pneumoniae⁺⁺⁺	73	100%		93%	100%	100%	97%	100%	100%	99%	97%	100%	99%	52%	97%	96%	(19)	95%
Proteus mirabilis	44	100%	85%	98%	100%	100%	98%	100%	100%	80%	89%		85%		93%	80%		
Pseudomonas aeruginosa	54	98%				89%		85%		89%	87%	98%	85%		98%		(32)	88%

Note: Isolates are counted once per patient (distinct count of medical record number + organism code).

+ CLS recommends that isolates reporting be restricted to counts >30. Interpret results with caution.

++ E. coli: 3.2% of isolates demonstrated ESBL.

+++ Klebsiella pneumoniae: 0% of isolates demonstrated ESBL.

*** Susceptibility for use in urinary tract infections, only.

**Sutter Medical Center Santa Rosa
Antibiogram 2012**

Organism Name	CountOfOrg code	Ampicillin	Cefazolin	Ceftriaxone	Ceftriaxone meningitis	Ciprofloxacin	Clindamycin	Erythromycin	Gentamicin	Levofloxacin	Linezolid	Nitrofurantoin++	Oxacillin	Pencillin	Tetracycline	Trimethoprim/Sulfa	Vancomycin+++
Enterococcus faecalis⁺	24	96%				92%		45%		92%	100%	100%			46%		96%
Enterococcus faecium⁺	14										100%						14%
Enterococcus species	54	94%				74%		100%		75%	100%	94%			24%		98%
Methicillin-resistant Staphylococcus aureus (MRSA)	100					36%	79%		99%	35%		100%			97%	94%	100%
Staphylococcus aureus	150		100%			85%	86%	68%	99%	86%		100%	100%		95%	97%	100%
Streptococcus pneumoniae⁺	6			100%	100%			80%						60%	50%	100%	100%

Note: Isolates are counted once per patient (distinct count of medical record number + organism code).

+ CLSI recommends that isolates reporting be restricted to counts >30. Interpret results with caution.

++ Susceptibility for use in urinary tract infections, only.